



MAG-FORCE

GROUP 2 FERTILIZER | Reg. No. B 6042 Act No. 36 of 1947

Mag-Force is a natural amino acid chelated plant nutrition containing magnesium for use as a foliar feed to maintain or increase magnesium levels in plants and fruits, for optimal crop growth and yield. Magnesium is essential for the formation of chlorophyll. It is also required for the metabolism of carbohydrates, plant cell respiration and plays a role in the synthesis of sugars, oils and fats within the plant. Mag-Force also contains macro- and micronutrients that is readily available.

COMPOSITION WHEN PACKED

Mg: 40 g/kg (45 g/ t)

SG @ 20°C: 1.26



MANUFACTURER:

Eco-Green Agriculture (Pty) Ltd

PO Box 3046, Rembrandt Mall, Paarl, 7620

Customer Care: +27 (0)21 871 1303

Fax: +27 (0)86 299 4198

info@ecogreenagri.com

www.ecogreenagri.com

Product Registration Holder: VIVRE



STIR & MIX THOROUGHLY BEFORE USE

Please see reverse side of label for information/direction for use



LIFT HERE

20 L

DIRECTION FOR USE

Use higher rates for crops cultivated under irrigation and for high plant density situations. Reduce rates proportionally when spraying smaller / younger plants and trees. The rate of application is also dependent on soil fertility and crop deficiencies. Lower rates should be applied for maintenance, and higher rates should be applied to correct nutritional deficiencies. Foliar applications should preferably be made during the cool times of the day. It is essential to obtain the specific recommendation from your representative before using this fertilizer.

CROP	FOLIAR APPLICATION RATE PER HA
Deciduous Fruit (Apples, Pears, Apricots, Peaches, Cherries, Fig, Nectarine, Plums, Pomegranates)	2 - 4 L Apply as a high volume foliar spray from fruit set. Apply 3 – 5 sprays throughout this growing period every 7 – 10 days.
Avocado Citrus Pecans Macadamia	2 - 4 L Apply as a high volume foliar spray during the growing season. Apply 3 – 5 sprays throughout this growing period every 7 – 10 days.
Grapes (Table & Wine)	2 - 4 L Apply as foliar spray two weeks after flowering. Repeat every 14 days.
Potatoes Onions	2 - 3 L First application can be made when deficiency symptoms appear. Repeat every 7 - 10 days.
Maize Soybeans Sunflower	2 - 3 L First application can be made when deficiency symptoms appear. Repeat every 7 - 10 days.
Wheat Oats Barley	2 - 3 L First application can be made when deficiency symptoms appear. Repeat every 7 - 10 days.
Vegetables	2 - 3 L First application can be made when deficiency symptoms appear. Apply 3 – 5 sprays as required.

MISCIBILITY:

Mag-Force is compatible with most agricultural remedies and nutrients. Do not mix Mag-Force with any copper containing fungicides, phosphate or phosphonate. Do not mix Mag-Force with highly alkaline material such as lime sulphur. If unsure about the miscibility and compatibility of the product, test mix before application. Where water volume differs from typical rates, adjust application rates accordingly.

Although this product has been extensively tested under a large variety of conditions, the manufacturer does not warrant that it will be efficacious under all conditions, because the action and effect thereof may be affected by factors such as abnormal soil, climatic and storage conditions, quality of dilution water, compatibility with other substances not indicated on the label as well as by the method, time and accuracy of application. The registration holder furthermore does not accept responsibility for damage to crops, vegetation, and the environment or harm to man or animal or for lack of performance of the product concerned due to failure of the user to follow label instructions or to the occurrence of conditions, which could not have been foreseen in terms of the registration. Consult the supplier in the event of any uncertainty.