NANO-TECHNOLOGY

CHLORIDE FREE

Approved by ECOCERT INPUTS
Controlled by ECOCERT SAS

Product suitable for use in Organic Farming in accordance with EU Regulations No 2018/848 and 2021/1165 and NOP Regulation.

GYP-FLO (SC)

GROUP 2 FERTILISER | Reg. No. B 5447 Act 36 of 1947



Gyp-Flo is a micronized liquid suspension concentrate of calcium and sulphate that can be applied through drip, micro or pivot irrigation systems and/or air, boomspray or any other agricultural crop-spraying equipment.

COMPOSITION WHEN PACKED

Ca: 122 g/kg (173 g/L)
S: 95 g/kg (135 g/L)
Fulvic Acid: 20 g/kg (28,4 g/L)
SG: 1,42 (± 0,03)
Particle size: D50 < 2 µm



For Marketing Purposes Only

eco-GREEN AGRICULTURE

MANUFACTURER:
Eco-Green Agriculture (Pty) Ltd
Suite 05, Private Bag 3041,
Northern Paarl, 7620, South Africa
Customer Care: +27 (0)21 871 1303
info@ecogreenagri.com
www.ecogreenagri.com

Product Registration Holder: VIVRE

Not classified as harzardous according to GHS.

20 L

STIR & MIX THOROUGHLY BEFORE USE

PLEASE SEE REVERSE SIDE OF LABEL FOR INFORMATION/DIRECTIONS FOR USE

KEEP OUT OF REACH OF CHILDREN AND ANIMALS

LIFT HERE

DIRECTIONS FOR USE: USE ONLY AS DIRECTED

IMPORTANT GUIDELINES:

- It is essential to obtain specific recommendations from the registration holder or a qualified agronomist before using this fertiliser. The rate will vary, taking into consideration soil type and analysis, stage of crop, crop deficiency, climatic conditions and time of year.

MIXING INSTRUCTIONS:

- The contents may settle, so always shake the container and stir the contents well before use.
- First, fill the tank halfway with water while the spray pump is circulating continuously.
- Slowly add the Gyp-Flo, then add the rest of the water to the tank.
- Always ensure that the solution is constantly agitated and circulated while applying.
- Applications through any type of irrigation system should always be made steadily and systematically, in other words, over the whole duration of the full irrigation cycle.

Use	Dosage Rate	Remarks
Row crops (e.g., wheat, canola, maize and potatoes)	**3 – 5 L/ha**	Apply in furrow next to seed row. Cal-Flo, Mag-Flo and Gyp-Flo may be applied together, depending on the soil analysis and recommendation made. During planting, it is advisable to apply Nutri-Soil in combination with Gyp-Flo for optimal plant growth and germination.
Drip/micro irrigation systems (e.g., vineyards, tree crops)	**10 – 20 L/ha**	Repeat as required. Allow 7 to 14 days between applications.
Sprinkler/centre pivot irrigation systems (e.g., potatoes, lucerne)	**10 – 20 L/ha**	Repeat as required. Allow 7 to 14 days between applications.
Greenhouse (e.g., vegetables, blueberries)	**1 – 2 ml Gyp-Flo per 1 L of planting medium (in bags)**	E.g., in a 14-litre bag, apply 14 to 28 ml of Gyp-Flo (depending on the sodium contents, Ca % etc.) Applications can be made by hand if necessary. Repeat as required. Allow 7 to 14 days between applications.

For Marketing Purposes Only

Although this product has been extensively tested under a large variety of conditions, the manufacturer does not warrant that it will be efficacious under all conditions, because the action and effect thereof may be affected by factors such as abnormal soil, climatic and storage conditions, quality of dilution water, compatibility with other substances not indicated on the label as well as by the method, time and accuracy of application. The registration holder furthermore does not accept responsibility for damage to crops, vegetation, and the environment or harm to man or animal or for lack of performance of the product concerned due to failure of the user to follow label instructions or to the occurrence of conditions, which could not have been foreseen in terms of the registration. Consult the supplier in the event of any uncertainty.